

Bibliographie

- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4e ed.). Washington, DC.
- Angst, J. (1990). Depression and anxiety: A review of studies in the community and in primary health care. Dans N. Sartorius, D. Goldberg, et al. (1990). *Psychological disorders in general medical settings* (pp. 60-68). Ashland, OH, US: Hogrefe & Huber Publishers.
- Appleby, L., Cooper, J., Amos, T., & Faragher, B. (1999). Psychological autopsy study of suicides by people aged under 35. *British Journal of Psychiatry*, 175(Aug), 168-174.
- Åsberg M (1986) Biochemical aspects of suicide. *Clinical Neuropharmacology*, 9(4), 374-376.
- Barraclough, B.M., Bunch, J., Nelson, B., & Sainsbury, P. (1974). A hundred cases of suicide: clinical aspects. *British Journal of Psychiatry*, 125, 355-373.
- Beautrais, A.L. (2003). Life course factors associated with suicidal behaviors in young people. *American Behavioral Scientist*, 46(9), 1137-1156.
- Beautrais, A.L., Joyce, P.R., Mulder, R.T., Fergusson, D.M., & Deavoll, B.J. (1996). Prevalence and comorbidity of mental disorders in persons making serious suicide attempts: a case-control study. *American Journal of Psychiatry*, 153(8), 1009-1014.
- Beck, A.T., Steer, R.A., Kovacs, M., Garrison, B. (1985). Hopelessness and eventual suicide: a 10-year prospective study of patients hospitalized with suicidal ideation. *American Journal of Psychiatry*, 142(5):559-63.
- Blair-West, G.W., Cantor, C.H., Mellsop, G.W., & Eyeson, A.M.L. (1999). Lifetime suicide risk in major depression: Sex and age determinants. *Journal of Affective Disorders*, 55(2-3), 171-178.
- Brent, D.A. (1988). Risk factors for adolescent suicide: A comparison of adolescent suicide victims with suicidal inpatients. *Archives of General Psychiatry*, 45(6), 581-588.
- Brent, D.A. (1994). Personality disorder, personality traits, impulsive violence and completed suicide in adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33(8), 1080-1086.
- Brent, D.A. (1995). Risk factors for adolescent suicide and suicidal behavior: Mental and substance abuse disorders, family environmental factors, and life stress. *Suicide and Life Threatening Behavior*, 25(Suppl), 52-63.
- Brent, D.A. (1999). Age- and sex-related risk factors for adolescent suicide. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38(12), 1497-1505.
- Brent, D.A., Perper, J.A., Moritz, G., Baugher, M., Roth, C., Balach, L., & Schweers, J. (1993). Stressful life events, psychopathology, and adolescent suicide: A case control study. *Suicide and Life Threatening Behavior*, 23(3), 179-187.
- Brown, S.L. (1985). Adolescents and family systems. In M. Peck, N.L. Farberow, & R.E. Litman (Eds.), *Youth suicide*. (pp. 71-79). New York: The Springer Series on Death and Suicide.

- Cavanagh, J.T., Carson, A.J., Sharpe, M., & Lawrie, S.M. (2003). Psychological autopsy studies of suicide : A systematic review. *Psychological Medicine*, 33(3), 395-405.
- Comité de la santé mentale du Québec (1994). *Recommandations pour développer et enrichir la Politique de santé mentale*, Comité de la santé mentale du Québec, Les Publications du Québec, Québec, 60 p.
- Conner, K.R. & Duberstein, P.R. (2004) Predisposing and precipitating factors for suicide among alcoholics : empirical review and conceptual integration. *Alcohol Clinical Experimental Research*, 28 (Suppl.5): 6S-17S.
- Cooper, J., Appleby, L., Amos, T. (2002). Life events preceding suicide by young people. *Social Psychiatry and Psychiatric Epidemiology*, 37(6):271-5.
- Dorpat, T., & Ripley, H. (1960). A study of suicide in the Seattle area. *Comprehensive Psychiatry*, 1, 349-359.
- Fenton, W.-S. (1997). Symptoms, subtype, and suicidality in patients with schizophrenia spectrum disorders. *American Journal of Psychiatry*, 154(2), 199-204
- Foster, T., Gillespie, K., & McClelland, R. (1997). Mental disorders and suicide in Northern Ireland. *British Journal of Psychiatry*, 170, 447-452.
- Gould, M.S. (1996). Psychosocial risk factors of child and adolescent completed suicide. *Archives of General Psychiatry*, 53, 1155-1162.
- Gould, M.S., Greenberg, T., Velting, D.M., & Shaffer, D. (2003). Youth suicide risk and preventive interventions: A review of the past 10 years. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42(4), 386-405.
- Groholt, B., Ekeberg, O., Wichstrom, L., & Haldorsen, T. (1998). Suicide among children and younger and older adolescents in Norway: A comparative study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37(5), 473-481.
- Hagnell, O., & Rorsman, B. (1979). Suicide in the Lundby study: a comparative investigation of clinical aspects. *Neuropsychobiology*, 5(2), 61-73
- Harris, E.C., & Barraclough, B.M. (1997). Suicide as an outcome for mental disorders. A meta-analysis. *British Journal of Psychiatry*, 170, 205-228.
- Heila, H., Isometsa, E.T., Henriksson, M.M., Heikkinen, M.E., Marttunen, M.J., & Lonnqvist, J.K. (1997). Suicide and schizophrenia: a nationwide psychological autopsy study on age- and sex-specific clinical characteristics of 92 suicide victims with schizophrenia. *American Journal of Psychiatry*, 154(9):1235-42.
- Houle, J. (2005). *La demande d'aide, le soutien social et le rôle masculin chez des hommes qui ont fait une tentative de suicide*. Thèse de doctorat. Université du Québec à Montréal.
- Houston, K., Hawton, K., & Shepperd, R. (2001). Suicide in young people aged 15-24: a psychological autopsy study. *Journal of Affective Disorders*, 63(1-3), 159-170.
- Institute of Medicine (1994). *Reducing risk for mental health disorders: Frontiers for preventive intervention research*. Washington, DC: National Academy Press.

- Kaplan, K.J., Harrow, M. (1996). Positive and negative symptoms as risk factors for later suicidal activity in schizophrenics versus depressives. *Suicide & Life Threatening Behaviour*, 26(2):105-21
- Kessler, R.C., McGonagle, K.A., Zhao, S., Nelson, C.B., Hughes, M., Eshleman, S., Wittchen, H.U., & Kendler, K.S. (1994). Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Archives of General Psychiatry*, 51(1), 8-19.
- Lesage, A.D., Boyer, R., Grunberg, F., Vanier, C., Morissette, R., Menard-Buteau, C., & Loyer, M. (1994). Suicide and mental disorders: A case-control study of young men. *American Journal of Psychiatry*, 151(7), 1063-1068.
- Lindstrom, M.B., Ryding, E., Bosson, P., Ahnide, J.A., Rosen, I., & Traskman-Bendz, L. (2004). Impulsivity related to brain serotonin transporter binding capacity in suicide attempters. *European Neuropsychopharmacology*, 14 (4) p.295-300.
- Malone, K.M., Haas, G.L., Sweeney, J.A., Mann, J.J. (1995). Major depression and the risk of attempted suicide. *Journal of Affective Disorders*,34(3):173-85
- Mann, J.J., Waternaux, C., Haas, G.L., & Malone, K.M. (1999). Toward a clinical model of suicidal behavior in psychiatric patients. *American Journal of Psychiatry*, 156(2):181-9.
- Marttunen, M.J., Aro, H.-M., Henriksson, M.M., & Lönnqvist, J.K. (1991). Mental disorders in adolescent suicide: DSM-III-R Axes I and II diagnoses in suicides among 13- to 19-year-olds in Finland. *Archives of General Psychiatry*, 48, 834-839.
- Meltzer, H.Y. (1999). Suicide and schizophrenia: clozapine and the InterSePT study. *International Clozaril/Leponex Suicide Prevention Trial. Journal of Clinical Psychiatry*. 60(Suppl.12):47-50.
- Michaelis, B.H., Goldberg, J.F., Davis, G.V., Singer, T.M., Garnon, J.L., & Wenzel, S.J. (2004). Dimensions of impulsivity and aggression associated with suicide attempts among bipolar patients: a preliminary study. *Suicide and Life Threatening Behavior*, 34(2): 172-176.
- Mishara, B.L., & Tousignant, M. (2004). *Comprendre le suicide*. Montréal: Presses de l'Université de Montréal.
- Nordstrom, P., Samuelsson, M., & Åsberg, M. (1994) CSF 5-HIAA predicts suicide risk after attempted suicide. *Suicide and Life Threatening Behavior*, 24, 1-9
- Organisation mondiale de la santé (2004). *Investir dans la santé mentale*. Produit par le Département de Santé mentale et abus de substances psychoactives - Maladies non transmissibles et santé mentale, Organisation mondiale de la Santé, Genève.
Accessible : http://www.who.int/mental_health/media/en/InvMHBr8.pdf
- Pfeffer, C.R. (1989). Life stress and family risk factors for youth fatal and nonfatal suicidal behavior. In C. R. Pfeffer (Ed.), *Suicide among youth: Perspectives on risk and prevention*. (pp. 143-164). Washington, D.C.
- Pirkis, J., & Burgess, P. (1998). Suicide and recency of health care contacts. A systematic review. *British Journal of Psychiatry*, 173, 462-474.

- Rich, C.L., Young, D., & Fowler, R.C. (1986). San Diego suicide study: I. Young vs old subjects. *Archives of General Psychiatry*, 43(6), 577-582.
- Robins, E. (1959). Some clinical considerations in the prevention of suicide based on a study of one hundred and thirty-four successful suicides. *American Journal of Public Health*, 49(7), 888-899.
- Robins, L.N. & Regier, D.A. (1991). *Psychiatric disorders in America: The Epidemiologic Catchment Area Study*. Free Press, New York.
- Rossau, C.D., & Mortensen, P.B. (1997). Risk factors for suicide in patients with schizophrenia: nested case-control study. *British Journal of Psychiatry*, 171(October), 335-339.
- Runeson, B.S., Beskow, J.E., & Waern, M. (1996). The suicidal process among young people. *Acta Psychiatrica Scandinavica*, 93(1), 35-42.
- Santé et Bien-être social Canada (1988). *La Santé mentale des Canadiens : vers un juste équilibre*. Ottawa.
- Shaffer, D., Gould, M.-S., Fisher, P., & Trautman, P., Moreau, D., Kleinman, M., & Flory, M. (1996). Psychiatric diagnosis in child and adolescent suicide. *Archives of General Psychiatry*, 53(4), 339-348.
- Shafii, M. (1988). Comorbidity of mental disorders in the post-mortem diagnosis of completed suicide in children and adolescents. *Journal of Affective Disorders*, 15(3), 227-233.
- Silverman, M.M., & Felner, R.D. (1995). Suicide prevention programs: Issues of design, implementation, feasibility, and developmental appropriateness. Special Issue: Suicide prevention: Toward the year 2000. *Suicide and Life Threatening Behavior*, 25(1), 92-104.
- Suominen, K., Henriksson, M., Suokas, J., Isometsa, E.T., Ostamo, A., & Lönnqvist, J.K. (1996). Mental disorders and comorbidity in attempted suicide. *Acta Psychiatrica Scandinavica*, 94(4), 234-240.
- Swan, A.C., Dougherty, D.M., Pazzaglia, P.J., Pham, M., & Moeller, F.G. (2004) Impulsivity: a link between bipolar disorder and substance abuse. *Bipolar Disorder*, 6(3):204-212.
- Tanney, B.L. (2000). Psychiatric diagnoses and suicidal acts. In R.W. Maris, A.L. Berman, & M.M. Silverman (Eds.), *Comprehensive Textbook of Suicidology*. (pp. 311-341). New York: The Guilford Press.
- U.S. Preventive Services Task Force. (1996). *Screening for Suicide Risk. Guide to Clinical Preventive Services* (pp.547-54), 2nd ed. Washington, DC: Office of Disease Prevention and Health Promotion.
- Ustun, T.B. & Sartorius, N. (1995). *Mental Illness in General Health Care: An International Study*. John Wiley & Sons.
- van Heeringen, C., Audenaert, K., Van Laere, K., Dumont, F., Slegers, G., Mertens, J., & Dierckx, R.A. (2003). Prefrontal 5-HT(2a) receptor binding index, hopelessness and personality characteristics in attempted suicide. *Journal of Affective Disorders*, 74(2), 149-58.